

---

# **AUSTRALIAN PRODUCT INFORMATION – KETAMINE INTERPHARMA 10 mg/mL and 50 mg/mL ampoules (KETAMINE HYDROCHLORIDE)**

## **1. NAME OF THE MEDICINE**

Ketamine hydrochloride

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

KETAMINE INTERPHARMA is formulated as an acid (pH 3.5 to 5.5) solution for intravenous or intramuscular injection in concentrations containing the equivalent of 10 mg or 50 mg ketamine base per millilitre.

For the full list of excipients, see Section 6.1 List of excipients.

## **3. PHARMACEUTICAL FORM**

KETAMINE INTERPHARMA is a clear, colourless solution for intravenous or intramuscular injection.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

KETAMINE INTERPHARMA is recommended:

1. as the sole anaesthetic agent for diagnostic and surgical procedures that do not require skeletal muscle relaxation. KETAMINE INTERPHARMA is best suited for short procedures and it can be used with additional doses, for longer procedures;
2. for the induction of anaesthesia prior to the administration of other general anaesthetic agents;
3. to supplement low-potency agents, such as nitrous oxide.

### **4.2 Dose and method of administration**

**Ketamine injection is for single use in one patient only. Discard any residue. All doses are given in terms of ketamine base.**

#### **Pre-Operative Preparation**

1. While vomiting has been reported following KETAMINE INTERPHARMA administration, airway protection is usually afforded because of active laryngeal-pharyngeal reflexes. However, because these reflexes may also be diminished by supplementary anaesthetics or muscle relaxants, the possibility of aspiration must be

---

considered. KETAMINE INTERPHARMA is recommended for use in the patient whose stomach is not empty only when, in the judgement of the medical practitioner, the benefits of the drug outweigh the possible risks.

2. Atropine, hyoscine or other 'drying' agents should be given at an appropriate interval prior to induction.

### **Dosage**

As with other general anaesthetic agents, the individual response to KETAMINE INTERPHARMA is somewhat varied depending on the dose, route of administration and age of patient, so that the dosage recommended cannot be absolutely determined in a fixed manner. The drug should be titrated against the patient's requirements.

### **Onset and Duration**

Because of rapid induction following the initial intravenous injection, the patient should be in a supported position during administration. The onset of action of KETAMINE INTERPHARMA is rapid; an intravenous dose of 2 mg/kg of body weight usually produces surgical anaesthesia within 30 seconds after injection, with the anaesthetic effect usually lasting 5 to 10 minutes. If a longer effect is desired, additional increments can be administered intravenously or intramuscularly to maintain anaesthesia without producing significant cumulative effect.

From experience, intramuscular doses (primarily in children, in a range of 9 to 13 mg/kg) usually produce surgical anaesthesia within 3 to 4 minutes following administration, with the anaesthetic effect usually lasting 12 to 25 minutes.

### **Induction**

**Intravenous route:** the initial dose of KETAMINE INTERPHARMA administered intravenously may range from 1 mg/kg to 4.5 mg/kg. The average amount required to produce 5 to 10 minutes of surgical anaesthesia has been 2 mg/kg.

### **NOTE**

**KETAMINE INTERPHARMA 10mg/mL and 50 mg/mL DO NOT REQUIRE DILUTION for intravenous injection.**

For dilution Ketamine Interpharma is compatible with normal saline or, 5% glucose in water.

**Rate of administration:** it is recommended that KETAMINE INTERPHARMA be administered slowly (over a period of 60 seconds). More rapid administration may result in respiratory depression and enhanced pressor response.

**Intramuscular route:** the initial dose of KETAMINE INTERPHARMA administered intramuscularly ranges from 6.5 to 13 mg/kg. A dose of 10 mg/kg will usually produce 12 to 25 minutes of surgical anaesthesia.

If the ketamine dose is augmented with diazepam, the two drugs must be given separately. Do **not** mix ketamine and diazepam in the same syringe or infusion flask.

---

**Dosage in Hepatic Insufficiency:** Dose reductions should be considered in patients with cirrhosis or other types of liver impairment (see Section 4.4 Special warnings and precautions for use).

### **Maintenance of Anaesthesia**

Increments of one half to the full induction dose may be repeated, as needed, for maintenance of anaesthesia. However it should be noted that involuntary and tonic-clonic movements of extremities might occur during the course of anaesthesia. These movements do not imply a level of attenuated anaesthesia and are not indicative of the need for additional doses of the anaesthetic. It should be recognised that the greater the total dose of KETAMINE INTERPHARMA administered, the longer will be the time to complete recovery.

This product is for one dose in one patient only. Discard any remaining contents.

### **4.3 Contraindications**

KETAMINE INTERPHARMA is contraindicated in patients with any condition in which a significant elevation of blood pressure would be hazardous such as: severe cardiovascular disease, heart failure, severe or poorly controlled hypertension, recent myocardial infarction, history of stroke, cerebral trauma, intracerebral mass or haemorrhage. Ketamine is also contraindicated in those who have shown hypersensitivity to the drug or its components.

### **4.4 Special warnings and precautions for use**

1. KETAMINE INTERPHARMA should be used by or under the direction of medical practitioners experienced in administering general anaesthetics and in maintenance of an airway and in the control of respiratory support.
2. Barbiturates and KETAMINE INTERPHARMA, being chemically incompatible because of precipitate formation, **should not** be injected from the same syringe.
3. Prolonged recovery time may occur if barbiturates and/or narcotics are used concurrently with KETAMINE INTERPHARMA.
4. Post-operative confusional states may occur during the recovery period (see Section 4.4 Special warnings and precautions for use - Emergence Reaction).
5. Because pharyngeal and laryngeal reflexes are usually active, KETAMINE INTERPHARMA should not be used alone in surgery or diagnostic procedures of the pharynx, larynx or bronchial tree. Mechanical stimulation of the pharynx should be avoided, whenever possible, if KETAMINE INTERPHARMA is used alone. Muscle relaxants with proper attention to respiration, may be required in both of these instances.
6. Resuscitative equipment should be ready for use.
7. The intravenous dose should be administered over a period of 60 seconds. More rapid administration may result in respiratory depression or apnoea and enhanced pressor response.

- 
8. In surgical procedures involving visceral pain pathways, KETAMINE INTERPHARMA should be supplemented with an agent, which obtunds visceral pain.
  9. Use with caution in the chronic alcoholic and the acutely alcohol-intoxicated patient. An increase in cerebrospinal fluid pressure has been reported following administration of KETAMINE INTERPHARMA. Use with extreme caution in patients with pre-anaesthetic elevated cerebrospinal fluid pressure.
  10. In patients with significant renal or hepatic impairment, the elimination of ketamine could potentially be delayed. Dose reductions should be considered in patients with cirrhosis or other types of liver impairment.
  11. Patients should be cautioned that driving an automobile, operating machinery or engaging in other hazardous activities should not be undertaken for 24 hours or more (depending on dose and other drugs employed) after anaesthesia.
  12. Use with caution in patients with increased intraocular pressure (eg. glaucoma) because the pressure may increase significantly after a single dose of ketamine
  13. Use with caution in patients with neurotic traits or psychiatric illness (eg. schizophrenia and acute psychosis)
  14. Use with caution in patients with acute intermittent porphyria
  15. Use with caution in patients with seizures
  16. Use with caution in patients with hyperthyroidism or patients receiving thyroid replacement (increased risk of hypertension and tachycardia)
  17. Use with caution in patients with pulmonary or upper respiratory infection (ketamine sensitises the gag reflex, potentially causing laryngospasm)
  18. Use with caution in patients with intracranial mass lesions, a presence of head injury, globe injuries, or hydrocephalus.

### **Emergence Reaction**

Treatment-emergent adverse reactions have occurred in approximately 12% of patients. The psychological manifestations vary in severity between pleasant dream-like states, vivid imagery, hallucinations, nightmares or illusions and delirium (often consisting of dissociative or floating sensations). In some cases, these states have been accompanied by confusion, excitement and irrational behaviour, which a few patients recall as an unpleasant experience. The duration ordinarily lasts no more than a few hours; in a few cases, however, recurrences have taken place up to 24 hours post-operatively. No residual psychological effects are known to have resulted from use of KETAMINE INTERPHARMA.

The incidence of these treatment-emergent adverse events is least in the young (15 years of age or less) and elderly (over 65 years of age) patient. Also they are less frequent when the drug is given intramuscularly. These reactions may be reduced if verbal, tactile and visual stimulation of the patient is minimised during the recovery period.

---

This does not preclude the monitoring of vital signs. In addition, the use of a small hypnotic dose of a short-acting or ultra-short-acting barbiturate may be required to terminate a severe treatment- emergent adverse reaction. The incidence of emergence reactions is reduced as experience with the drug is gained. When KETAMINE INTERPHARMA is used on an out-patient basis, the patient should not be released until recovery of anaesthesia is complete and should be accompanied by a responsible adult at discharge.

### **Cardiovascular**

Because of the substantial increase in myocardial oxygen consumption, ketamine should be used with caution in patients with hypovolemia, dehydration, or cardiac disease, especially coronary artery disease (eg. congestive heart failure, myocardial ischaemia, and myocardial infarction). In addition ketamine should be used with caution in patients with mild-to- moderate hypertension and tachyarrhythmias.

Cardiac function should be continually monitored during the procedure in patients found to have hypertension or cardiac decompensation.

### **Abuse Potential**

Ketamine has been reported being used as a drug of abuse. Reports suggest that ketamine produces a variety of symptoms including, but not limited to, flashbacks, hallucinations, dysphoria, anxiety, insomnia, or disorientation. Ketamine dependence and tolerance may develop in individuals with a history of drug abuse or dependence. Therefore, ketamine should be prescribed and administered with caution.

### **Use in the elderly**

No data available.

### **Paediatric use**

Plasma half-life, clearance and volume of distribution (relative to body weight) are not significantly different between adults and children, although absorption following intramuscular injection is more rapid in the latter.<sup>1, 2</sup>

### **Paediatric Neurotoxicity**

Some published studies in children have observed cognitive deficits after repeated or prolonged exposures to anaesthetic agents early in life. These studies have substantial limitations, and it is not clear if the observed effects are due to the anaesthetic/analgesic/sedation drug administration or other factors such as the surgery or underlying illness.

Published animal studies of some anaesthetic/analgesic/sedation drugs have reported adverse effects on brain development in early life and late pregnancy. The clinical significance of these nonclinical finding is yet to be determined

With inhalation or infusion of such drugs, exposure is longer than the period of inhalation or infusion. Depending on the drug and patient characteristics, as well as dosage, the elimination phase may be prolonged relative to the period of administration.

### **Effects on laboratory tests**

---

There is no information available regarding the possible effects of ketamine on clinical laboratory tests.

#### 4.5 Interactions with other medicines and other forms of interactions

Halogenated hydrocarbon inhalational anaesthetics may prolong the half-life of ketamine; recovery from anaesthesia may be prolonged following concurrent use. Concurrent use of ketamine (especially in high doses or when rapidly administered) with halogenated anaesthetics can increase the risk of developing bradycardia, hypotension, or decreased cardiac output.

Diazepam is known to increase the half-life of ketamine and prolongs its pharmacodynamic effects. Dose adjustments may therefore be needed<sup>3</sup>.

Sympathomimetics (directly or indirectly acting) and vasopressin may enhance the sympathomimetic effects of ketamine<sup>3</sup>.

Prolonged recovery time may occur if barbiturates and/or narcotics are used concurrently with ketamine.

Benzodiazepines may prolong the half life of ketamine; recovery from anaesthesia may be prolonged following concurrent use.<sup>4</sup>

Concomitant use with ergometrine may lead to an increase in blood pressure and Co-administration of drugs with a hypertensive effect should be avoided.<sup>3,5</sup>

Sustained rises in arterial pressure have been reported in patients receiving concomitant ketamine and thyroxine.<sup>5</sup>

Clinically significant reduction in seizure threshold may be observed in patients receiving concomitant ketamine and theophylline or aminophylline.<sup>3,5</sup> Unpredictable extensor-type seizures have been reported with concurrent administration of these agents.

There is no information available on the interactions between ketamine and antihypertensive agents. However, given the marked increase in arterial pressure following administration of ketamine, cardiac function should be monitored (see Section 4.4 Special warnings and precautions for use).

Barbiturates and KETAMINE INTERPHARMA, being chemically incompatible because of precipitate formation, **should not** be injected from the same syringe.

Ketamine is clinically compatible with the commonly used general and local anaesthetic agents when an adequate respiratory exchange is maintained.

Ketamine may potentiate the neuromuscular blocking effects of atracurium and tubocurarine, including respiratory depression with apnoea.

The use of ketamine with other central nervous system (CNS) depressants (eg. ethanol, phenothiazines, sedating H<sub>1</sub>-blockers, or skeletal muscle relaxants) can potentiate CNS depression and/or increase risk of developing respiratory depression. Reduced doses of ketamine may be required with concurrent administration of other anxiolytics, sedatives, and

---

hypnotics.

Ketamine has been reported to antagonise the hypnotic effect of thiopental.

Patients taking thyroid hormones have an increased risk of developing hypertension and tachycardia when given ketamine.

Concomitant use of antihypertensive agents and ketamine increases the risk of developing hypotension.

## **4.6 Fertility, pregnancy and lactation**

### **Effects on fertility**

No data available.

### **Use in pregnancy**

Category B3

Limited studies in animals have not shown that ketamine causes birth defects; however, it crosses the placenta. Histological changes in the heart (degeneration and oedema of cardiac muscle), liver (diffuse haemopoietic cell infiltration, parenchymal cell degeneration) and kidneys (proximal convoluted tubule degeneration) were observed in foetuses following administration of ketamine to pregnant rats during the period of organogenesis at doses similar to the maximum human dose, on a body surface area basis; a NOEL for these effects was not established. Ketamine administration to pregnant monkeys near term was associated with increased blood pCO<sub>2</sub> and a dose-dependent respiratory depression in neonates, at a dose about one sixteenth the maximum human dose on a body surface area basis.

With the exception of administration during surgery for abdominal delivery or vaginal delivery, no controlled clinical studies in pregnancy have been conducted. The safe use of ketamine in pregnancy has not been established, and such use is not recommended.

Australian categorisation definition of Category B3:

Drugs which have been taken by only a limited number of pregnant women and women of childbearing age, without an increase in the frequency of malformation or other direct or indirect harmful effects on the human fetus having been observed.

Studies in animals have shown evidence of an increased occurrence of fetal damage, the significance of which is considered uncertain in humans.

Published animal studies of some anaesthetic/analgesic/sedation drugs have reported adverse effects on brain development in early life and late pregnancy

Published studies in pregnant and juvenile animals demonstrate that the use of anaesthetic/analgesic and sedation drugs that block NMDA receptors and/or potentiate GABA activity during the period of rapid brain growth or synaptogenesis may result in neuronal and oligodendrocyte cell loss in the developing brain and alterations in synaptic morphology and neurogenesis when used for longer than 3 hours. These studies included anaesthetic agents from a variety of drug classes.

---

## **Use in lactation**

Ketamine is likely to be excreted in breast milk and therefore breastfeeding should be discontinued when ketamine is in use.

## **4.7 Effects on ability to drive and use machines**

Patients should be cautioned that driving an automobile, operating machinery or engaging in other hazardous activities should not be undertaken for 24 hours or more (depending on dose and other drugs employed) after anaesthesia.

## **4.8 Adverse effects (undesirable effects)**

### **Cardiovascular**

Blood pressure and pulse rate are frequently elevated following administration of KETAMINE INTERPHARMA. However, hypotension and bradycardia have been observed. Arrhythmia has also occurred.

### **Respiration**

Although respiration is frequently stimulated, severe depression of respiration or apnoea may occur following rapid intravenous administration of high doses of KETAMINE INTERPHARMA. Laryngospasm and other forms of airway obstruction have occurred during KETAMINE INTERPHARMA anaesthesia.

### **Eye**

Diplopia and nystagmus have been noted following KETAMINE INTERPHARMA administration. KETAMINE INTERPHARMA may also cause a slight elevation in intraocular pressure measurement.

### **Psychological (see Section 4.4 Special warnings and precautions for use - Emergence Reaction)**

### **Neurological**

In some patients, enhanced skeletal muscle tone may be manifested by tonic and clonic movements, sometimes resembling seizures (see Section 4.2 Dose and method of administration).

### **Gastrointestinal**

Anorexia, nausea and vomiting have been observed. However this is not usually severe and allows the great majority of patients to take liquids by mouth shortly after regaining consciousness (see Section 4.2 Dose and method of administration). Hypersalivation has also been observed.

### **Abuse Potential (see Section 4.4 Special warnings and precautions for use)**

---

## Immune System Disorders

Anaphylaxis has been observed.

## General

Local pain and exanthema at the injection site have infrequently been reported. Transient erythema and/or morbilliform rash have also been reported.

## Reporting suspected adverse effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

## 4.9 Overdose

Respiratory depression may occur with overdosage or too rapid rate of administration of KETAMINE INTERPHARMA, in which case, supportive ventilation should be employed. Mechanical support of respiration is preferred to administration of analeptics.

Ketamine has a wide margin of safety; several instances of unintentional administration of overdoses of KETAMINE INTERPHARMA (up to 10 times that usually required) have been followed by prolonged but complete recovery.

For information on the management of overdose, contact the Poisons Information Centre on 13 11 26 (Australia).

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

#### Mechanism of action

KETAMINE INTERPHARMA is a rapid-acting, general anaesthetic producing an anaesthetic state characterised by profound analgesia, normal pharyngeal-laryngeal reflexes, normal or slightly enhanced skeletal muscle tone, cardiovascular and respiratory stimulation, and occasionally, a transient and minimal respiratory depression.

A patent airway is maintained, partly by virtue of relatively unimpaired pharyngeal and laryngeal reflexes (see **Section 4.4 Special warnings and precautions for use**).

The anaesthetic state produced by KETAMINE INTERPHARMA has been termed 'dissociative anaesthesia' in that it appears to selectively interrupt association pathways of the brain before producing somesthetic sensory blockade. KETAMINE INTERPHARMA may selectively depress the thalamocortical system before significantly obtunding the more ancient cerebral centres and pathways (reticular-activating and limbic systems).

---

Elevation of blood pressure begins shortly after injection, reaches a maximum within a few minutes and usually returns to pre-anaesthetic values within 15 minutes after injection. The median peak rise has ranged from 20 to 25% of pre-anaesthetic values.

### **Clinical trials**

KETAMINE INTERPHARMA ketamine (as hydrochloride) has been studied in over 12,000 operative and diagnostic procedures involving over 10,000 patients from 105 separate studies. During the course of these studies, KETAMINE INTERPHARMA was administered as the sole agent, as induction for other general anaesthetic agents, or to supplement low potency agents. In these studies, the anaesthesia was rated either “excellent” or “good” by the anaesthetist and the surgeon at 90% and 93% respectively. In a second method of evaluation, the anaesthesia was rated “adequate” in at least 90% and “inadequate” in 10% or less of procedures. Specific areas of application have included the following:

1. debridement, painful dressings and skin grafting in burn patients as well as other superficial surgical procedures;
2. neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures;
3. diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions;
4. diagnostic and operative procedures of the pharynx, larynx or bronchial tree;

Note: muscle relaxants with proper attention to respiration, may be required (see **Section 4.4 Special warnings and precautions for use**)

5. sigmoidoscopy and minor surgery of the anus and rectum and circumcision;
6. extraperitoneal procedures used in gynaecology, such as dilation and curettage;
7. orthopaedic procedures such as closed reductions, manipulations, femoral pinning, amputations and biopsies;
8. as an anaesthetic in poor-risk patients with depression of vital functions;
9. in procedures where the intramuscular route of administration is preferred;
10. in cardiac catheterisation procedures.

## **5.2 Pharmacokinetic properties**

### **Absorption**

Ketamine is rapidly absorbed following parenteral administration. Peak plasma levels averaged 0.75µg/ml and CSF levels were about 0.2µg/ml one hour after dosing.<sup>6</sup> The plasma half-life is in the range of 2 to 4 hours.<sup>7,8,9</sup> After IM administration (absorption half-life 2-17 minutes) it is up to 93 % bioavailable.<sup>6</sup>

---

## Distribution

Ketamine (as hydrochloride) is rapidly and extensively distributed throughout the body into highly perfused tissues including the brain.<sup>8,9</sup> Mean volume of distribution is reported to range from approximately 1 to 3 L/kg, and the distribution half-life is approximately 7 to 11 minutes. Ketamine (as hydrochloride) is approximately 20-50% bound to plasma proteins.<sup>4</sup> Ketamine is likely to be excreted in breast milk, but this is unlikely to be clinically relevant. The drug crosses the placenta in induction doses<sup>10</sup> (see Section 4.6 Fertility, pregnancy and lactation).

## Metabolism<sup>4</sup>

Ketamine undergoes extensive hepatic metabolism. The biotransformation includes N-dealkylation to norketamine (metabolite I), hydroxylation of the cyclohexone ring (metabolites III and IV), conjugation with glucuronic acid and dehydration of the hydroxylated metabolites to form the cyclohexene derivative (metabolite II). Norketamine (metabolite I) has about 1/6 of the potency of ketamine and is formed at concentrations in the plasma similar to those of the parent compound.

## Excretion

After intravenous bolus administration, ketamine shows a bi- or triexponential pattern of elimination. The alpha phase lasts about 45 minutes with a half-life of 10 to 15 minutes. This first phase, which represents the anaesthetic action of ketamine, is terminated by redistribution from the CNS to peripheral tissues and hepatic biotransformation to an active metabolite. The beta phase half-life is about 2.5 hours.<sup>7,8,9</sup> About 90% of ketamine is excreted in the urine, mostly as metabolites, with only about 2 to 4 % as the unchanged drug. Approximately 5% is recovered in the faeces.<sup>11</sup> The renal clearance of ketamine hydrochloride is  $15 \pm 5$  mL/min/kg.<sup>12</sup>

## 5.3 Preclinical safety data

### Genotoxicity

No data available.

### Carcinogenicity

No data available.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Water for injections

10mg/mL preparations have saline added to achieve isotonicity.

### 6.2 Incompatibilities

Barbiturates and KETAMINE INTERPHARMA, being chemically incompatible because of

---

precipitate formation, should not be injected from the same syringe.

## 6.2 Shelf life

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

## 6.3 Special precautions for storage

Store below 30°C. Protect from light.

KETAMINE INTERPHARMA should not be used if the solution is coloured and/or contains particulate matter.

## 6.4 Nature and contents of container

### **KETAMINE INTERPHARMA 10 mg/mL presented in the following sizes:**

KETAMINE INTERPHARMA 20 mg/2 mL solution for injection (AUST R 310918)

KETAMINE INTERPHARMA 50 mg/5 mL solution for injection (AUST R 310910)

KETAMINE INTERPHARMA 100 mg/10mL solution for injection (AUST R 310913)

### **KETAMINE INTERPHARMA 50 mg/mL presented in the following sizes:**

KETAMINE INTERPHARMA 100 mg/2 mL solution for injection (AUST R 310914)

KETAMINE INTERPHARMA 250 mg/5 mL solution for injection (AUST R 310912)

KETAMINE INTERPHARMA 500 mg/10mL solution for injection (AUST R 310916)


Type I clear glass ampoules in packs of 5 ampoules  
Not all strengths may be marketed.

## 6.5 Special precautions for disposal

In Australia, any unused medicine or waste material should be disposed of in accordance with local requirements.

## 6.6 Physicochemical properties

### Chemical structure



---

ketamine (as hydrochloride) is a non-barbiturate anaesthetic chemically designated (2*RS*)-2-(2-Chlorophenyl)-2-(methylamino) cyclohexanone hydrochloride. Ketamine is a racemic mixture. The molecular weight is 274.2 and the empirical formula is C<sub>13</sub>H<sub>16</sub>ClNO.HCl.

Ketamine is freely soluble in water and methyl alcohol and is soluble in alcohol.

**CAS number**

1867-66-9

**7. MEDICINE SCHEDULE (POISONS STANDARD)**

Schedule 8 - Controlled Drug

**8. SPONSOR**

InterPharma Pty Ltd  
Suite 103,  
39 East Esplanade  
Manly, NSW, 2095  
Australia  
Tel: +61 29976 6876  
Web: www.interpharma.com.au

**9. DATE OF FIRST APPROVAL**

16 December 2019

**10. DATE OF REVISION**

**Summary Table of Changes**

<b>Section changed</b>	<b>Summary of new information</b>

- 
1. Nimmo WS, et al. Pharmacokinetics of ketamine in children. Br J Anaesth 1982; 14: 144P.
  2. Grant IS, et al. Ketamine disposition in adults and children Br J Anaesth 1983, 55: 1107-11.2.5 Clinical Overview Updates to Section 4.5 Interaction With Other Medicinal Products and Other Forms of Interaction of the Core Data Sheet November 2016.
  3. Therapeutic Drugs. Edited by Sir Colin Dollery, 1991, Vol 2: K7-13.
  4. Martindale The Complete Drug Reference, MICROMEDEX Healthcare series Vol. 105 IncCopyright 2000 Pharmaceutical Press.

- 
5. Clements JA, Nimmo WS, Grant IS. Bioavailability, pharmacokinetics and analgesic activity of ketamine in humans. *J Pharm Sci* 1982; 71: 539-41.
  6. Clements JA, Nimmo WS. Pharmacokinetics and analgesic effects of ketamine in man. *Br J Anaesth* 1981; 53: 27-30.
  7. Grant IS, et al. Pharmacokinetics and analgesic affects of IM and oral ketamine. *Br. J Anaesth* 1981; 53: 805-9.
  8. Wieber J, Gryler RD, Hengstmann JH, Dengler HJ. Pharmacokinetics of ketamine in man. *Anaesthesist* 1975; 24: 260-6.
  9. Little B, Chang T, Chaucet L, et al. A study of ketamine as an obstetrical anesthetic. *Am J Obstet Gynecol* 1972; 113: 247-58.
  10. United States Pharmacopeia Dispensing Information, 1998, 18<sup>th</sup> Edition, pg 1775-7.
  11. Geisslinger G, et al. Pharmacokinetics and pharmacodynamics of ketamine enantiomers in surgical patients using a stereoselective analytical method. *Br J Anaesth* 1993, 70: 666-71.